

Physical Risks

Life-Threatening Risks of Abortion

Higher death risk, up to 7 times higher suicide

Compared to pregnant women who had their babies, pregnant women who aborted were ...

- **3.5 times more likely to die in the following year**
- **1.6 times more likely to die of natural causes**
- **6-7 times more likely to die of suicide**
- **14 times more likely to die from homicide**
- **4 times more likely to die of injuries related to accidents¹**

Another study found that, compared to women who gave birth, women who had abortions had a 62% higher risk of death from all causes for at least *eight* years after their pregnancies. Deaths from suicides and accidents were most prominent, with deaths from suicides being 2.5 times higher.²

Causes of death within a week — The leading causes of abortion-related maternal deaths within a week of abortion are hemorrhage, infection, embolism, anesthesia complications, and undiagnosed ectopic pregnancies.³

Cancer — Significantly increased risk of breast cancer, cervical cancer, and lung cancer (probably due to heavier smoking patterns after abortion).⁴

Immediate complications — About 10% suffer immediate complications; one-fifth are life-threatening:⁵

- hemorrhage
- cervical injury
- convulsions
- endotoxic shock
- infection
- embolism
- chronic abdominal pain
- second-degree burns
- ripped or perforated uterus
- anesthesia complications
- gastro-intestinal disturbances
- Rh sensitization

31% suffer health complications— A recent study published in a major medical journal found that 31% of American women surveyed who had undergone abortions had health complications.⁶

80%-180% increase in doctor visits — Based on health care sought before and after abortion. On average, there is an 80% increase in doctor visits and a 180% increase in doctor visits for psychosocial reasons after abortion.⁷

Self-destructive lifestyles, spiraling health problems — Increased risk of promiscuity, smoking, drug abuse, and eating disorders, which all put the woman at increased risk for other health problems.⁸

Infertility and life-threatening reproductive risks

Abortion can damage reproductive organs and cause long-term and sometimes permanent problems that can put future pregnancies at risk. Women who have abortions are more likely to experience ectopic pregnancies, infertility, hysterectomies, stillbirths, miscarriages, and premature births than women who have not had abortions.⁹

Teens Face Higher Risk, 6 Times More Likely to Attempt Suicide

Teens 6 times more likely to attempt suicide— Teenage girls are 10 times more likely to attempt suicide if they have had an abortion in the last 6 months than are teens who have not had an abortion.¹⁰

Reproductive damage and other complications — Compared to teens who give birth, teens who abort are generally at higher risk of immediate complications and long-term reproductive damage after abortion than are older women.¹¹

continued ►

Higher risk of PID, 2.5 times higher risk of endometritis (a major cause of maternal death in future pregnancies)

— Teens are at higher risk for dangerous infections such as pelvic inflammatory disease and endometritis after abortion. These infections increase their risk of infertility, hysterectomy, ectopic pregnancy, and other serious complications.¹²

Overview of reproductive complications and problems with subsequent deliveries

Pelvic Inflammatory Disease — Abortion puts women at risk of Pelvic Inflammatory Disease (PID) is a serious, life-threatening disease and a major direct cause of infertility. PID also increases risk of ectopic pregnancies. Studies have found that approximately one-fourth of women who have a chlamydia infection at the time of their abortion and 5% of women who don't have chlamydia will develop PID within four weeks after the abortion.¹³

Placenta Previa — After abortion, there is a seven- to 15-fold increase in placenta previa in subsequent pregnancies (a life-threatening condition for both the mother and her wanted pregnancy). Abnormal development of the placenta due to uterine damage increases the risk of birth defects, stillbirth, and excessive bleeding during labor.¹⁴

Ectopic Pregnancy — Post-abortive women have a significantly increased risk of subsequent ectopic pregnancies,¹⁵ which are life threatening and may result in reduced fertility.

Endometritis, a Major Cause of Death — Abortion can result in for endometritis, which can lead to hospitalization and infertility problems. It is a major cause of maternal death during pregnancy.¹⁶

Women who abort twice as likely to have pre-term or post-term deliveries.¹⁷

- Women who had one, two, or more previous induced abortions are, respectively, 1.89, 2.66, or 2.03 times more likely to have a subsequent pre-term delivery, compared to women who carry to term. Pre-term delivery increases the risk of neonatal death and handicaps. The average hospital charge from delivery to discharge for a premature birth is \$58,000, compared to \$4,300 for a full-term birth.
- Women who had one, two, or more induced abortions are, respectively, 1.89, 2.61, and 2.23 times more likely to have a post-term delivery (over 42 weeks).

Death or disability of newborns in later pregnancies — Cervical and uterine damage may increase the risk of premature delivery, complications of labor, and abnormal development of the placenta in later pregnancies.¹⁶ These complications are the leading causes of disabilities among newborns.

AfterAbortion.org/news

The Elliot Institute was involved in all of the studies listed above. For more information on this research, including citations and links to the published studies, visit www.afterabortion.org/news.

Detrimental Effects -- Quick-Reference Summary of Available Research

Additional research summaries are available in the book “Detrimental Effects of Abortion: An Annotated Bibliography with Commentary,” edited by Thomas W. Strahan. To order, call **Acorn Books: 1-888-412-2676**.

Citations for *Physical Risks of Abortion* Fact Sheet

1. M Gissler et. al., “Pregnancy Associated Deaths in Finland 1987-1994 -- definition problems and benefits of record linkage,” *Acta Obstetrica et Gynecologica Scandinavica* 76:651-657, 1997; Mika Gissler, Elina Hemminki, Jouko Lonnqvist, “Suicides after pregnancy in Finland: 1987-94: register linkage study” *British Medical Journal* 313:1431-4, 1996; and M. Gissler, “Injury deaths, suicides and homicides associated with pregnancy, Finland 1987-2000,” *European J. Public Health* 15(5):459-63, 2005 .
2. DC Reardon et. al., “Deaths Associated With Pregnancy Outcome: A Record Linkage Study of Low Income Women,” *Southern Medical Journal* 95(8):834-41, Aug. 2002.

3. Kaunitz, "Causes of Maternal Mortality in the United States, *Obstetrics and Gynecology* 65(5), May 1985
4. H.L. Howe, et al., "Early Abortion and Breast Cancer Risk Among Women Under Age 40," *International Journal of Epidemiology* 18(2):300-304, 1989; L.I. Remennick, "Induced Abortion as A Cancer Risk Factor: A Review of Epidemiological Evidence," *Journal of Epidemiological Community Health* 1990; M.C. Pike, "Oral Contraceptive Use and Early Abortion as Risk Factors for Breast Cancer in Young Women," *British Journal of Cancer* 43:72, 1981; M-G, Le, et al., "Oral Contraceptive Use and Breast or Cervical Cancer: Preliminary Results of a French Case-Control Study, *Hormones and Sexual Factors in Human Cancer Etiology* ed. JP Wolff, et al., (New York, Excerpta Medica, 1984) 139-147; F. Parazzini, et al., "Reproductive Factors and the Risk of Invasive and Intraepithelial Cervical Neoplasia," *British Journal of Cancer* 59:805-809, 1989; H.L. Stewart, et al., "Epidemiology of Cancers of the Uterine Cervix and Corpus, Breast and Ovary in Israel and New York City," *Journal of the National Cancer Institute* 37(1):1-96; I. Fujimoto, et al., "Epidemiologic Study of Carcinoma in Situ of the Cervix," *Journal of Reproductive Medicine* 30(7):535, July 1985; N. Weiss, "Events of Reproductive Life and the Incidence of Epithelial Ovarian Cancer," *Am. J. of Epidemiology*, 117(2):128-139, 1983; V. Beral, et al., "Does Pregnancy Protect Against Ovarian Cancer," *The Lancet* 1083-7, May 20, 1978; C. LaVecchia, et al., "Reproductive Factors and the Risk of Hepatocellular Carcinoma in Women," *International Journal of Cancer* 52:351, 1992.
5. Frank, et.al., "Induced Abortion Operations and Their Early Sequelae," *Journal of the Royal College of General Practitioners* 35(73):175-180, April 1985; Grimes and Cates, "Abortion: Methods and Complications", in *Human Reproduction*, 2nd ed., 796-813; M.A. Freedman, "Comparison of complication rates in first trimester abortions performed by physician assistants and physicians," *Am. J. Public Health* 76(5):550-554, 1986).
6. VM Rue et. al., "Induced abortion and traumatic stress: A preliminary comparison of American and Russian women," *Medical Science Monitor* 10(10): SR5-16, 2004.
7. P. Ney, et.al., "The Effects of Pregnancy Loss on Women's Health," *Soc. Sci. Med.* 48(9):1193-1200, 1994; Badgley, Caron, & Powell, *Report of the Committee on the Abortion Law* (Ottawa: Supply and Services, 1997) 319-321.
8. T. Burke with D. Reardon, *Forbidden Grief: The Unspoken Pain of Abortion* (Springfield, IL: Acorn Books, 2002), see ch. 13 and 15.
9. Strahan, T. *Detrimental Effects of Abortion: An Annotated Bibliography with Commentary* (Springfield, IL: Acorn Books, 2002) 168-206.
10. B. Garfinkle, *Stress, Depression and Suicide: A Study of Adolescents in Minnesota* (Minneapolis: University of Minnesota Extension Service, 1986).
11. Wadhwa, "Legal Abortion Among Teens, 1974-1978", *Canadian Medical Association Journal* 122:1386-1389 June 1980; 13. E. Belanger, et. al., "Pain of First Trimester Abortion: A Study of Psychosocial and Medical Predictors," *Pain*, 36:339; G.M. Smith, et. al., "Pain of first-trimester abortion: Its quantification and relationships with other variables," *American Journal Obstetrics & Gynecology*, 133:489, 1979; R.T. Burkman, et. al., "Morbidity Risk Among Young Adolescents Undergoing Elective Abortion," *Contraception*, 30(2):99, 1984; and K.F. Schulz, et. al., and "Measures to Prevent Cervical Injury During Suction Curettage Abortion," *The Lancet*, 1182-1184, May 28, 1993 .
12. Burkman, et al., "Morbidity Risk Among Young Adolescents Undergoing Elective Abortion" *Contraception* 30:99-105, 1984; R.T. Burkman, et. al., "Culture and treatment results in endometritis following elective abortion," *American J. Obstet. & Gynecol.*, 128:556, 1997; and D. Avonts and P. Piot, "Genital infections in women undergoing induced abortion," *European J. Obstet. & Gynecol. & Reproductive Biology*, 20:53, 1985; W. Cates, Jr., "Teenagers and Sexual Risk-Taking: The Best of Times and the Worst of Times," *Journal of Adolescent Health*, 12:84, 1991; and "Teenage Pregnancy: Overall Trends and State-by-State Information," Report by the Alan Guttmacher Institute, Washington, DC, www.agi.org.
13. Radberg, et al., "Chlamydia Trachomatis in Relation to Infections Following First Trimester Abortions," *Acta Obstetrica Gynaecologica* (Supp. 93), 54:478, 1980; L. Westergaard, "Significance of Cervical Chlamydia Trachomatis Infection in Post-abortal Pelvic Inflammatory Disease," *Obstetrics and Gynecology* 60(3):322-325, 1982; M. Chacko, et al., "Chlamydia Trachomatis Infection in Sexually Active Adolescents: Prevalence and Risk Factors," *Pediatrics* 73(6), 1984; M. Barbacci, et al., "Post-Abortal Endometritis and Isolation of Chlamydia Trachomatis," *Obstetrics and Gynecology* 68(5):668-690, 1986; S. Duthrie, et al., "Morbidity After Termination of Pregnancy in First-Trimester," *Genitourinary Medicine* 63(3):182-187, 1987.
14. Barrett, et al., "Induced Abortion: A Risk Factor for Placenta Previa", *American Journal of Ob&Gyn.* 141:7, 1981.
15. Daling, et.al., "Ectopic Pregnancy in Relation to Previous Induced Abortion", *J. American Medical Association* 253(7):1005-1008, Feb. 15, 1985; Levin, et.al., "Ectopic Pregnancy and Prior Induced Abortion", *American J. Public Health* 72:253, 1982; C.S. Chung, "Induced Abortion and Ectopic Pregnancy in Subsequent Pregnancies," *American J. Epidemiology* 115(6):879-887 (1982).
16. "Post-Abortal Endometritis and Isolation of Chlamydia Trachomatis," *Obstetrics and Gynecology* 68(5):668- 690, 1986); P. Sykes, "Complications of termination of pregnancy: a retrospective study of admissions to Christchurch Women's Hospital, 1989 and 1990," *New Zealand Medical Journal* 106: 83-85, March 10, 1993; S Osser and K Persson, "Postabortal pelvic infection associated with Chlamydia trachomatis infection and the influence of humoral immunity," *Am J Obstet Gynecol* 150:699, 1984; B. Hamark and L. Forssman, "Postabortal Endometritis in Chlamydia-Negative Women- Association with Preoperative Clinical Signs of Infection," *Gynecol Obstet Invest* 31:102-105, 1991; and Strahan, *Detrimental Effects of Abortion: An Annotated Bibliography With Commentary* (Springfield, IL: Acorn Books, 2002) 169.
17. Zhou, Weijin, et. al., "Induced Abortion and Subsequent Pregnancy Duration," *Obstetrics & Gynecology* 94(6):948-953, Dec. 1999.
18. Hogue, Cates and Tietze, "Impact of Vacuum Aspiration Abortion on Future Childbearing: A Review", *Family Planning Perspectives* 15(3), May-June 1983.