60% said, "part of me died."

- 1. 64% of women having abortions felt pressured by others.<sup>1</sup> Coercion can escalate to violence or even murder.<sup>1, 2</sup>
- 2. 52% felt rushed and 54% were not sure about the decision at the time, yet an astounding 67% received *no* counseling beforehand.<sup>1</sup>
- 3. 84% reported that they did not receive adequate counseling.<sup>1</sup>
- 4. 79% were not informed about available alternatives.<sup>1</sup>
- 5. 31% of women suffered health complications after abortion.<sup>1</sup> 10% suffer immediate complications, one-fifth of which are life-threatening.<sup>3</sup>
- 6. Women have a 65% higher risk of clinical depression after abortion compared to women who give birth.<sup>4</sup>
- 7. 65% suffer multiple symptoms of post-traumatic stress disorder after abortion.<sup>1</sup>
- 8. Death rates from all causes are 3.5 times higher among women who abort, compared to women who give birth.<sup>5</sup>
- 9. 60% said "part of me died," according to a survey of women who aborted.<sup>1</sup>
- 10. Suicide rates are 6 times higher compared to women who give birth.<sup>6</sup>

## Citations

1. VM Rue et. al., "Induced abortion and traumatic stress: A preliminary comparison of American and Russian women," *Medical Science Monitor* 10(10): SR5-16 (2004).

2. See the special report, "Forced Abortion in America," at www.theunchoice.com/resources.htm.

3. Frank, et.al., "Induced Abortion Operations and Their Early Sequelae," *Journal of the Royal College of General Practitioners* 35(73):175-180 (April 1985); Grimes and Cates, "Abortion: Methods and Complications", in *Human Reproduction*, 2nd ed., 796-813; and M.A. Freedman, "Comparison of complication rates in first trimester abortions performed by physician assistants and physicians," *Am. J. Public Health* 76(5):550-554 (1986).

4. JR Cougle, DC Reardon & PK Coleman, "Depression Associated With Abortion and Childbirth: A Long-Term Analysis of the NLSY Cohort," *Medical Science Monitor* 9(4): CR105-112 (2003).

5. M Gissler et. al., "Pregnancy Associated Deaths in Finland 1987-1994 -- definition problems and benefits of record linkage," Acta Obsetricia et Gynecologica Scandinavica 76:651-657 (1997).

6. Mika Gissler, Elina Hemminki, Jouko Lonnqvist, "Suicides after pregnancy in Finland: 1987-94: register linkage study," *British Medical Journal* 313:1431-4, 1996; and M. Gissler, "Injury deaths, suicides and homicides associated with pregnancy, Finland 1987-2000," *European J. Public Health* 15(5):459-63 (2005).